

Biodiversity Climate change impacts

report card technical paper

4. Implications of Climate Change for SSSIs and other Protected Areas

Philippa Gillingham

School of Applied Sciences
Bournemouth University
Fern Barrow
Poole
Dorset
BH12 5BB

01202 962372

Email: pgillingham@bournemouth.ac.uk

Executive Summary

Sites of Special Scientific Interest (SSSIs) cover just under 7 % of England, around 12 % of Wales and around 13 % of Scotland. In recent years, investments have been made to bring the habitats and species that SSSIs were designated for into 'favourable' condition. However, some SSSIs and other protected areas (PAs) are under direct physical threat from inundation following sea level rise, and changes in climate will affect the species and habitats that are present on most PAs. This report summarises these threats and considers options for changing the way that the protected area network is managed.

The following impacts of climate change have already been detected on PAs;

- Saltmarshes have been lost to coastal squeeze, and coastal freshwater habitats including grazing marsh and lowland raised bog are at risk of inundation by seawater under current conditions
- Northern species have decreased in density, whilst southern species have increased in density. Whilst most evidence for this occurring on PAs comes from outside the UK, there is evidence that these patterns are occurring within the UK as well
- Southern species in the UK have used PAs to facilitate their northwards expansion and many occur in higher numbers on PAs compared to other areas in the newly colonised parts of their range
- Northern species that have experienced a decrease in range have persisted more often in PAs than elsewhere.

In addition, the following future impacts of climate change can be anticipated;

- The composition of flora and fauna on each PA will change – high confidence (**medium evidence, high agreement**)
- Cold adapted species of high latitudes and altitudes will tend to decrease on PAs, whilst warm adapted species will tend to increase – medium confidence (**medium evidence, medium agreement**)
- PAs in the North of the UK will gain plant species overall, whilst PAs in the south may lose some native plant species. This pattern is reversed for UK breeding birds – low confidence (**medium evidence, low agreement**)
- Species with lower dispersal capacities and those for which urban and intensive agricultural areas are a barrier to dispersal will be unable to colonize PAs that become climatically suitable – low confidence (**limited evidence, medium agreement**)
- Work in Africa predicted that some Important Bird Areas (IBAs) may lose all the species for which they were designated by 2085, although for around 90 % of species at least one currently occupied IBA should remain suitable, similarly in Asia median predicted turnover by 2100 was 43 %, but no species were predicted to lose all climate space across the network. In Europe, species turnover is predicted to be faster than in Africa – medium confidence (**medium evidence, medium agreement**)
- Increasing range mismatching of interacting species, such as butterflies and their host plants, might mean that more management is necessary on PAs to

preserve species that interact with each other – low confidence (**limited evidence, medium agreement**)

- Hotspots of bird diversity in Finland and Norway may no longer coincide geographically with PA boundaries – low confidence (**limited evidence, medium agreement**)

Integrating consideration of climate change into management plans for the PA network is likely to result in more effective (and cost-effective) conservation solutions. In order to facilitate this integration, monitoring of climate change impacts and management actions should be carried out to enable adaptive decision making.

Introduction

Protected areas (PAs) cover 10.1-15.5 % of the globe (depending on definition; Chape *et al.* 2005, Soutullo 2010), and the Aichi Targets of the Convention on Biological Diversity aim to increase this to 17% (Harrop 2011). In the UK, there are many different conservation designations. Areas designated primarily for the protection of biodiversity cover just under 7 % of England, around 12 % of Wales and around 13 % of Scotland, (Sites of Special Scientific Interest (SSSI) which encompass Special Areas of Conservation (SACs, designated under the EU Habitats Directive), Special Protection Areas (SPAs, designated under the EU Birds Directive) and Ramsar sites (designated under the Ramsar convention)). The area protected increases to over 23 % of England and over 25 % of Wales if all designations are considered, including those primarily designated for the landscape character, such as Areas of Outstanding Natural Beauty and National Parks (Lawton *et al.* 2010). Even these less biodiversity-focussed designations may decrease the likelihood of activities that are potentially harmful to wildlife occurring (Bate *et al.* 2010).

Patterns in the distribution and size of the UK's protected sites

Protected areas are not spread uniformly across the UK, so this paper begins by examining patterns in the distribution and size of the UK's protected sites. Ancient Woodland habitats are covered by the planning system as well as by SSSI designation (less than 22 % of broadleaved woodland in England is within SSSIs, Lawton *et al.* 2010), so patterns in their distribution have been investigated in addition to SSSIs. Within the UK, 1 km grid squares at higher altitudes tend to have a higher percentage cover of SSSI (Spearman's Rank Correlation, $\rho=0.28$, $P<0.0001$), reflecting the greater proportion of semi-natural habitat in the uplands. Conversely, 1 km grid squares with high percentage cover of ancient woodland are often found at lower altitudes (Spearman's Rank Correlation, $\rho=0.05$, $P<0.0001$ – the relationship is very weak despite the highly significant p-value). In addition to this, across the whole of the UK, higher latitudes tend to have a higher percentage of SSSI (Spearman's Rank Correlation, $\rho=0.26$, $P<0.0001$), whilst the distribution of Ancient Woodland shows a bimodal distribution with latitude, so that there is a higher percentage of cover at low and high latitudes, with less at the central latitudes. This pattern in distribution reflects the land use history of different soil types and topographies. Many areas of ancient woodland in the UK are very small (modal value is 0.29 hectares) although this differs between countries – England and Wales have many small areas of ancient woodland (mode 0.28 hectares for England, 0.27 hectares for Wales, >44,000 recorded sites in England, >48,000 recorded sites in

Wales), whilst Scotland has fewer but larger areas of this habitat (mode 1.97 hectares, <29,000 recorded sites).

Importance of Protected Areas to conservation

In the UK, where areas outside PAs are often highly modified, some plant species are entirely confined to PAs and most are well represented in PAs, although some critically endangered species appear not to be represented within their borders (Jackson *et al.* 2009). Across Europe, there is some evidence that SPAs improve the population trends of the species they were designed to protect. Donald *et al.* (2007) found an association between the percentage of land protected and the population trend of European breeding birds between two survey periods. This relationship was stronger for species protected under Annex 1 of the Birds Directive, for which SPAs are designated, but still held for non-Annex 1 species. A similar pattern was found in Eastern Europe (Koleček *et al.* 2014), whereby population trends of protected birds improved more than those that were not protected.

However, the contribution of SSSIs to conserving non-target species can be variable. In a study of eight British butterflies, Davies *et al.* (2007) found that whilst population trends tended to be positive on SSSIs, half the species studied maintained higher populations on SSSIs in unfavourable condition than they did on SSSIs in favourable condition according to common standards monitoring. They concluded that management for biodiversity in Britain is detrimental to butterflies associated with later seral stages of grassland and scrub/grassland mosaics, which suggests there may be more that can be done to conserve biodiversity on SSSIs. A more recent study (Brereton *et al.* 2011) found that population trends for 12 specialist butterflies were no different on SSSIs than on unprotected land, and numbers of these species were declining across the UK.

In addition to benefitting biodiversity, SSSIs also contribute significant benefits in terms of the ecosystem services they provide. Christie and Rayment (2012) found that the general public in England and Wales were willing to pay almost nine times more than the current cost of SSSI management for the ecosystem services they provide. In addition, it appears that designated sites may deliver more cultural and regulating services than non-designated sites (Eastwood *et al.* 2013) although this study involved relatively few sites.

Vulnerability of Protected Areas to Climate Change

Protected areas have different vulnerabilities to the effects of climate change depending on their location, size, sensitivity of component habitats and species, current condition and the presence of non-climatic factors such as pollution (Wilson *et al.* 2010). Coastal habitats are particularly vulnerable to inundation by seawater and coastal erosion. For example, large areas of many saltmarshes protected in SPAs have been lost to coastal squeeze since their designation (Haskoning 2006). Fresh water habitats located close to the coast are also at risk from sea-level rise (DEFRA 2011) with 1,531 ha of SSSI habitat in England (3.5 % of total SSSI area present in the coastal floodplain, including grazing marsh and lowland raised bog) at risk under current (2010) conditions. This increases to 4.7 % by 2100 under a medium emissions scenario with degraded defences, and recovery from inundation is not guaranteed. Recent wetland creation work has resulted in an increase in the number of sites occupied by Bittern *Botaurus stellaris*. However, this population

growth could be threatened by the loss of just three sites in Suffolk that are currently under threat from sea level rise (Gilbert *et al.* 2010).

Small PAs are less likely to retain areas with similar climatic conditions in the future (Loarie *et al.* 2009) so are less likely to retain the species that are currently resident than larger PAs due to a lack of climate connectivity (Hodgson *et al.* 2009). Because important habitats tend to be more fragmented in England and Wales than Scotland (see section on distribution of PAs), we might expect a higher level of vulnerability in these two countries.

Different taxonomic groups may also have different vulnerabilities. In work carried out by the Joint Nature Conservation Committee, fish were the species group least likely to have been assessed as favourable, and lowland and upland heath were the habitat types with the lowest percentages assessed as favourable under common standards monitoring (Williams 2006). Since current condition can affect vulnerability to climate change, these habitats might therefore be expected to be especially vulnerable. In more recent work (Brooker *et al.* 2014), features in Scotland were ranked by their vulnerability to climate change (based on expert opinion). The majority of the highest risk features were assemblages of species. This was followed by freshwater and wetland habitats. In England, coastal and wetland Special Protection Areas (SPAs) are at higher risk than other SPAs (Franks *et al.* 2014).

Implications of changing species distributions for Protected Areas

A concern for some authors is that PAs are fairly static in space, whilst species respond to climate change by moving their distributions. This potential problem was recognised as early as 1985, when Peters and Darling (1985) used paleoecological data to show that the predominant response of species to climate change was to shift their distributions to more suitable locations. Despite this early recognition of the problem, as recently as 2004, climate change was not recognised globally by reserve managers as a potential threat to conserving species within PAs (WWF 2004). Climate change is expected to become a particular problem at the southern range margins of species distributions, where species may move out of reserves which were designated for them (Araújo *et al.* 2004, 2011). In recent years, some reserves have been degazetted in response to loss of the species that they were designated for (Mascia and Pailler 2011), and this could become a problem in the UK, particularly in reserves designated for one or a few species or habitat types.

So far only one study has specifically assessed the current effectiveness of protected areas in retaining UK species with retracting ranges (Gillingham *et al.* 2015a). Using data from repeat surveys of four northern butterflies and six northern birds, they concluded that there was a noticeable effect of protection on the likelihood of persistence of species, but that this positive effect of protection was lower at higher latitudes and altitudes. Studies such as this are very difficult to do for most species due to the requirement for repeat surveys of the same locations to determine where extinctions have occurred within the UK. In Australia, there have been calls to replace 'underperforming' PAs (Fuller *et al.* 2010). Other authors have also discussed the possibility of declassifying and selling some reserves in order to purchase or designate others (Strange *et al.* 2011). However, the performance criteria used by Fuller *et al.* (2010) did not include the potential future effectiveness

of PAs under changed climatic conditions. In the UK, where many PAs are privately owned, degazetting one PA would not free up funds for the designation of others.

Because many species have expanded their distributions northwards into new areas in response to climate change (e.g. Hickling *et al.* 2006), we should expect that species will disperse into PAs as well as out of them. In support of this theory, there is evidence that a wide range of invertebrate species, as well as some birds, disproportionately colonise SSSIs in Great Britain when expanding their distributions northwards (Thomas *et al.* 2012) and for those with more detailed data available, this was not simply because PAs tend to be situated at higher altitudes and latitudes in Great Britain (Gillingham *et al.* 2015a). In addition, many butterflies and odonates occur in higher numbers in PAs compared to non-designated areas in their newly colonised range (Gillingham *et al.* 2015b) despite these PAs not being designated or officially managed for them. Hiley *et al.* (2013) also found evidence that birds naturally colonising the UK from continental Europe have used PAs as 'landing pads' to facilitate this colonisation, yet PAs did not facilitate the spread of species that were introduced by humans (Hiley *et al.* 2014). This means that PAs should continue to be useful locations for conserving biodiversity in the UK in future, even if some species move out of them. Specialists were found to be more reliant on SSSIs than generalists (Thomas *et al.* 2012), suggesting that it is the habitats found within UK PAs that drive this pattern. However, this latter effect was only investigated for colonising butterflies, so the extent to which this rule holds for other taxa is a knowledge gap at present.

Reserve managers in the UK already monitor and manage habitats for some species that reserves were not originally designated for (Davies *et al.* 2007), and this presents an opportunity to increase the biodiversity under protection in the UK, which is one of the few countries in Europe predicted to 'win' overall in terms of the numbers of species that should find climatic conditions suitable in the future (IPCC 2007), with only higher latitude and altitude areas of Europe gaining species by 2020, 2050 and 2080 (IPCC 2014). This sort of pattern has been picked up in other European countries; on PAs in Finland, northern bird species have decreased in density in recent years, whilst southern species have increased, probably in response to a changing climate (Kujala *et al.* 2011, Virkkala and Rajasärkkä 2011). Despite decreases in richness of birds across Finland, PAs remain the places with highest richness and trends have been less negative on PAs for birds preferring mires (Virkkala *et al.* 2014). These studies, whilst not in the UK, give confidence in the general effectiveness of PAs as a conservation strategy under climate change.

1. Lessons from modelling bird distributions in sub-saharan Africa, Europe and Asia

Using climate envelope models, Hole *et al.* (2009) modelled the potential future distributions of 1,608 bird species breeding in 803 Important Bird Areas (IBAs). 815 of these were 'priority' species for which IBAs are designated. For 88-92 % of priority species (depending on climate scenario used) at least one of the IBAs projected to be climatically suitable in 2085 is currently designated for the species (i.e. there is an overlap in current and projected future range), and less than 1 % are projected to lose all suitable climate space within the network by 2085. However, 51-55 % of IBAs are projected to lose all the priority species for which they are currently designated, and range extent for priority species is projected to decline to 74 % of the current area occupied. In addition, in parts of the continent IBAs are separated by distances of > 500 km, which is substantially greater than the dispersal distances even bird species are capable of attaining, especially if the intervening terrain is inhospitable.

Modelling the distributions of 487 breeding birds in Europe, Huntley *et al.* (2010) found that species turnover was predicted to be higher and persistence lower in PAs in Europe than in Africa, with the 156 Annex 1 species projected to be particularly affected. However, northern Scotland was predicted to have high persistence of species, probably reflecting the high topographic heterogeneity and hence wide range of microclimatic conditions present.

In Asia, none of the modelled 370 species were projected to lose all suitable climate space from the IBA network by 2100 despite declines in the suitability of climate for 45% of species (Bagchi *et al.* 2013). However, there was a median turnover of 43% on individual IBAs.

Lessons learned – Reserve networks can be effective in protecting biodiversity in the short term even without significant dispersal of species, and PAs in the UK can contribute to international conservation objectives. However, in the longer term, actions that help species move between protected areas (such as habitat creation between isolated reserves, sympathetic management of areas surrounding reserves or even assisted colonisation) may be important. Flexibility in reserve designation may also be required, since some reserves may well lose all the species they have been designated for. These conclusions are echoed in part by a study from North America that looked at trees, birds, mammals and amphibians (Lawler and Hepinstall-Cymerman 2010).

Projecting future impacts of climate change on PAs

Modelling work in the US has shown that the composition of flora and fauna on individual PAs is likely to change (Lawler and Hepinstall-Cymerman 2010), and other studies have predicted that the representation of northern biomes on PAs will decrease, whilst representation of southern biomes will increase in both Canada and the UK (Lemieux and Scott 2005, Trivedi *et al.* 2008). Some PAs in Africa may lose all the species for which they were designated (Hole *et al.* 2009, Huntley *et al.* 2012, see box 1), and the likelihood of this occurring will increase with more severe climatic change. Despite these losses, around 90 % of the bird species modelled were predicted to retain suitable climatic conditions in at least one PA. In addition,

colonisation of PAs by expanding bird species will mean that some PAs will likely gain species overall, whilst others will lose species overall.

PAs in the north of the UK have been predicted to gain plant species, whilst PAs in the south are likely to lose plant species (Dockerty *et al.* 2003, see box 3). This pattern is reversed for UK breeding birds however (Pearce-Higgins *et al.* 2011, see box 4), so there is uncertainty in what the relative impacts of climate change will be on different taxonomic groups and sites. Using predicted range changes of species, PAs can be classified as likely to have high persistence of species, increasing specialisation, high predicted turnover, increasing value or increasing diversification. These different types of PA will have different optimum management strategies (Hole *et al.* 2011) if their biodiversity is to be conserved effectively.

<p>2. UK Bryophytes and Climate Change</p> <p>Anderson and Ohlemüller (2011) modelled the distributions of 43 rare UK bryophytes under current and future climates. Across all species, there was an increase in coverage within protected areas of the area considered climatically suitable, from 8.9 % in 1990-2020 to 10.2 % in 2051-2080. Suitable climate space moves uphill and to higher latitudes, which have a higher percentage covered within the protected area network. However, the median overlap between the current range and the climate space that is predicted to be suitable in future decreases from 21 % in 1990-2020 to 10 % in 2051-2080. At least a quarter of all species have no overlap between their current distribution and areas with analogous climates in future. Many species may therefore find it difficult to migrate to areas that are suitable in the future.</p>	<p>3. Plants in UK protected areas</p> <p>Dockerty <i>et al.</i> (2003) classified 200 species as 'declining', 'increasing' or 'no change' between current and projected future distributions on 66 nature reserves in the UK. Future moisture levels and temperature were projected to stay within the ranges already experienced by all 200 species in some part of their current European ranges. Northerly sites were projected to have more increasing trends in the probability of occurrence, whilst southerly sites were predicted to have more decreasing trends. At a single reserve (Backwarden SSSI in Essex), depending on the climate scenario used, 17-20 % of species showed increasing probability of occurrence, 28-48 % showed no change and 29-49 % showed a decreasing trend, including 3 species identified as of conservation priority by the site managers. Warming is likely to favour southern-temperate and Mediterranean types, with arctic-montane and boreal-montane types likely to decline (Trivedi <i>et al.</i> 2008).</p>
--	---

Not all species will be able to colonise all newly suitable PAs (see boxes 1 and 2). Species with lower dispersal capacities and those that are sensitive to urban barriers, such as the pool frog, may find it difficult to colonize areas that become climatically suitable (BRANCH partnership 2007, Anderson and Ohlemüller 2011, Pellatt *et al.* 2012). Because species have different dispersal capabilities, there may in future be an increasing mismatch in the ranges of interacting species, such as

butterflies and their host plants (Schweiger *et al.* 2012), although a lack of data on dispersal distances and how this interacts with habitat fragmentation to affect the ability of species to track climate change (Hodgson *et al.* 2012) mean that it is difficult to assess where these mismatches might occur. Projections of bird distributions in Finland and Norway showed that in those countries, even if species colonise new areas, hotspots of species diversity may no longer coincide geographically with PA boundaries (Virkkala *et al.* 2010), although PAs should remain useful for birds preferring mires, marshes and arctic mountains (Virkkala *et al.* 2013). In the UK, where there is extensive human-dominated use of the landscape between PAs, this is less likely since PAs represent the most suitable places for many species to colonise (Thomas *et al.* 2012).

4. The impact of climate change on UK birds on SPAs (CHAINSPAN project)

An important criterion for SPA designation is based on numerical thresholds – for instance, whether >1% of the UK population of a species is present at a site, or whether the site holds an important congregation of species. Changes in abundance at individual sites could therefore cause SPAs to be degazetted if they become less suitable for the species they were selected for. These criteria also result in poor representation of certain species groups, such as migratory passerines, in the UK SPA network.

Until recently, most modelling of potential climate impacts looked at impacts on range, rather than the population criterion on which designation is based. Pearce-Higgins *et al.* (2011) therefore projected future changes in abundance of bird species at individual Special Protection Areas (SPAs) as a result of climate change, using data from the UK, Ireland, the Netherlands and France. Sufficient data existed to fit 118 models (including separate models for some species in two seasons). Climate was a reasonable predictor of distribution across the models fitted, but it was a weak predictor of abundance at individual sites. In 33 of the models, climate had very low predictive power and only six models fitted the data well ($r > 0.5$), with climate accounting for approximately 19 % of variation in recent population trends across all populations modelled. This suggests that other factors are currently more important in determining the abundances of birds within individual SPAs. Therefore, site-based management is likely to be of use in reducing the adverse effects of climate change. Many species were projected to respond favourably to climate change in the short term, but with increasing severity of change, a greater proportion of species were projected to decline. The most vulnerable groups were predicted to be northern breeding seabirds and terrestrial species. Northern SPAs were predicted to lose qualifying features whilst many southern sites were predicted to gain features, but larger sites should continue to support more birds.

Several knowledge gaps were highlighted, as there were insufficient data to produce models for several of the most threatened terrestrial species. There is currently no requirement to take climate change into account when creating future management plans. An investigation of potential management responses on SPAs has now followed up the CHAINSPAN project, focussed on 4 contrasting SPAs (Franks *et al.* in press). Managers of these reserves identified a range of measures that would help them to best manage these sites taking into account climate change, including increased flexibility of agri-environment schemes and SPA designation processes and investment in water infrastructure. Separately the IPENS (Improvement programme for England's Natura 2000 sites) project (Natural England, 2015) has developed an approach to climate change adaptation across SPAs (and SACs) in England.

Clearly, sites designated for several species or habitats of interest will be less vulnerable to being degazetted as species move in response to climate change than sites designated for one or a few species or habitats. Since the presence of Annex-listed species determines the management of Natura 2000 sites (Verschuuren 2010), adding species of conservation concern to designation criteria as they colonise new areas should result in a lower likelihood of degazettement along with appropriate management being planned to ensure long-term survival of these colonising species. However, species with their warm range margins within the UK, for which dispersal to new areas would be difficult, could be disadvantaged by such actions if they were to result in less available suitable habitat. A complementary strategy would be to integrate expected shifts in species' distributions due to climate change into the conservation strategy of the Habitats Directive (as suggested by Normand *et al.* 2007).

Managing climate change impacts in PAs

Climate change adaptation has become well accepted policy and practise for nature conservation in the UK and elsewhere; it is for example set out in the National Adaptation Programme (HM Government 2013) and Biodiversity 2020 strategy (Defra 2011b). Natural England and the RSPB have recently (2014) published a climate change adaptation manual to support conservation practitioners in adaptation.

Reserve managers include adaptation to climate change in their plans and consider the wider reserve network (Franks *et al.* in press, MacGregor and van Dijk 2014). In some cases, appropriate management may have potential to reduce the negative impact of climate change by reducing the impacts of other negative drivers of population density (Pyke and Marty 2005, Pearce-Higgins 2011a, Singh and Milner-Gulland 2011). For example, the Golden Plover *Pluvialis apricaria* is dependent on the abundance of adult craneflies (Diptera: Tipulidae), which are a key prey item during the breeding season (Pearce-Higgins *et al.* 2010). Re-wetting peat by blocking drainage channels is expected to increase the numbers of craneflies available as a food source (Carroll *et al.* 2011), which may help the Golden Plover persist in the face of climate change. However, there are trade-offs to be considered with such a course of action. A single management action can affect species of conservation concern differentially. For example, intensively managed grouse moors in upland Britain are associated with lower declines of Lapwing *Vanellus vanellus* but faster declines of Golden Plover *P. apricaria* (Amar *et al.* 2011). In addition, management to restore one vegetation type can negatively impact the cover of other desirable plants (Mitchell *et al.* 2009).

Another potential management action is to create habitat suitable for expanding species in areas that become climatically suitable for them. The Royal Society for the Protection of Birds (RSPB) has recently re-created heathland suitable for the Dartford Warbler from a conifer plantation, to enable it to colonise new areas during its current northwards expansion (RSPB 2010). There is also recent evidence that PAs under active management have been colonised more frequently by the Silver spotted skipper butterfly *Hesperia comma* (Lawson *et al.* 2014). Habitat management may also be necessary if translocations are to be considered to aid species to track climate change (e.g. Willis *et al.* 2009). There is a trade-off to consider here, as maintaining habitat for retreating species may discourage


expanding species from colonising protected areas. In a world where conservation resources are limited, some management actions may use resources that could potentially be spent elsewhere. In the UK, beech *Fagus sylvatica* has sometimes been removed from woodland in the North West, where it is was not previously found, whereas in Southern England, where conditions are becoming less suitable for it, it has been managed to enable persistence (Gaston *et al.* 2006). In more recent years this position has been revised in light of climate change to be more accepting of beech colonising northern and western areas (Natural England 2009). Where species distributions are considered in terms of whole range dynamics, these changes in management actions should allow the species in question to colonise areas that have become climatically suitable (Monzón *et al.* 2011). Conservation management in the UK is increasingly recognising this need to consider site scale conservation within the wider context of species ranges (e.g. Natural England and RSPB 2014).

Facilitating movement across the landscape could result in more cost-effective conservation outcomes, enabling resources to be redirected. This would require new interest features to be added to reserve management plans, whilst species that have been irreversibly lost from a site would be removed from the designation and management objectives to ensure efficient use of resources, and this approach has been identified as desirable by managers of SPAs at risk from climate change (Franks *et al.* in press). In all cases, new objectives for individual PAs should take into account species' wider conservation status and distributions (Dodd *et al.* 2012). This approach is already adopted by Natural England (Natural England 2012), who committed to revise the conservation objectives for SSSIs and develop new SSSI guidance that will take into account climate change issues. They will also implement a Notification Strategy which includes a boundary and feature review of all SSSIs, ensuring that climate change adaptation is considered. Macgregor and van Dijk (2014) interviewed nature reserve managers in Eastern England to assess whether and how climate change was being incorporated into their management plans. A variety of strategies were being used, from attempting to maintain current species or ecosystems, to facilitating movement of species and helping new species establish (MacGregor and van Dijk 2014). Monitoring of the success of these different strategies will be instrumental in informing conservation practise into the future.

Habitat associations of species change along climatic gradients (Oliver 2009, Suggitt *et al.* 2012), which may complicate the picture further as it will be difficult for reserve managers to predict how best to manage for a particular species in the future. At the leading edge of species' ranges, habitat breadth can expand as the climate becomes more suitable, enabling species to exploit a wider range of habitats (Pateman *et al.* 2012), which might facilitate range expansions in response to climate change. Given this information, it is reasonable to expect that habitat breadth might decrease at the trailing edge of species' ranges, as the climate becomes less suitable. In addition, other environmental drivers can reduce habitat breadth despite climatic release (Oliver *et al.* 2012). Hen Harrier *Circus cyaneus* shows differential responses to management at different sites within the UK, meaning that habitat management guidelines have had to be developed on a site by site basis (Arroyo *et al.* 2005). The likely changes of habitat preference under climatic change and at different latitudes and elevations is a major knowledge gap, as most of the available literature is concentrated on the Lepidoptera, and it is not currently known to what extent other species might follow similar patterns.

Adapting the PA network to ameliorate climate change

Based on the best current knowledge, Lawton *et al.* (2010) concluded that England's protected area network would need to be modified in order to adapt to the challenges posed by climate change. Several options have been proposed when designating new PAs in response to climate change. Because climate change and habitat fragmentation act synergistically to decrease the abundance and range of species (Opdam and Wascher 2004), many authors have suggested increasing physical connectivity between habitats (Heller and Zavaleta 2009) or temporal connectivity between suitable climate space (Hodgson *et al.* 2009) by including areas of topographic and climatic heterogeneity within PAs (Carroll *et al.* 2010). Some authors advocate designating dynamic PAs to complement existing static ones in the marine environment (Game *et al.* 2009), but this would be difficult to achieve within the UK's highly modified terrestrial environment. The use of dynamic reserves are constrained by habitat fragmentation outside reserves, necessitating management of the matrix (Rayfield *et al.* 2008), and other authors have stressed that expanding and connecting reserve networks will be insufficient to conserve biodiversity under climate change (Kostyack *et al.* 2011), so management of land between reserves will be necessary anyway. Others have suggested that reserves should be designated based on criteria that include future performance (Singh and Milner-Gulland 2011) or that new reserves should be established in the expected direction of travel of suitable climate space (Pearson and Dawson 2005). These approaches are species-centric and could be expensive to apply to a large number of species, many of which will have competing demands, as well as relying on uncertain model predictions. However, only a small amount of additional land may be necessary to create a climatically robust representation of some species (Pyke and Fischer 2005), and if this did not involve a high economic cost might be worth considering when designating new reserves. Monzón *et al.* (2011) suggested that management should be changed within reserves to take account of the dynamism of species responses to climate change, so that resources are not wasted on maintaining species at a site once the climate has become unsuitable, assuming that the species in question has expanded its distribution elsewhere. The use of long-term monitoring of population densities will be important in detecting initial responses to climate change, as well as the effectiveness of any management actions (see Figure 1).



Based on the information provided in this report, the UK Biodiversity Partnership adaptation principles (Hopkins *et al.* 2007), and the England Biodiversity Strategy (Smithers *et al.* 2008) several recommendations can be made (for a more general discussion of adaptive management for climate change consult Natural England and RSPB 2014);

1. Existing PAs should be retained. PAs protect large percentages of most important semi-natural habitats, without this protection the land might be used for activities harmful to biodiversity (Lawton *et al.* 2010). Reducing fragmentation and increasing patch size may make some species more resilient to climate change (Newson *et al.* 2014).
2. Management within PAs could help to reduce sources of harm not linked to climate change, for example (Pearce-Higgins 2011a) by decreasing predation rates and increasing available prey resources for birds. The effectiveness of these management interventions in decreasing vulnerability to climate change should be monitored and conservation priorities regularly reviewed to ensure resources are used efficiently (e.g. Pearce-Higgins 2011a, 2011b, see figure 1).
3. Maintaining heterogeneity within landscapes should increase the chances that species will be able to spread locally into newly favourable habitat (Hodgson *et al.* 2009, Carroll *et al.* 2010). Heterogeneity within sites is associated with dampened population dynamics (i.e. reducing the chance of stochastic events causing extinction, Oliver *et al.* 2014) and in any case both vegetation and topographic heterogeneity appear to be drivers of species richness for a wide range of taxa (Stein *et al.* 2014).
4. Creating new habitat (Hodgson *et al.* 2011), restoring degraded habitat, or reducing the intensity of management of the landscape between existing habitats

- should facilitate species' movements between PAs. The RSPB aims to double the area it currently manages for nature conservation by 2030 (RSPB 2007).
5. When reviewing management plans, the likely future impacts of climate change should be considered and appropriate changes made (Monzón *et al.* 2011, Pearce-Higgins 2011b, see figure 1). This approach has been taken by the RSPB in their futurescapes campaign (Dodd *et al.* 2010, RSPB 2010).
 6. Intertidal habitat should be re-created and protected through managed realignment, to compensate for losses predicted by coastal squeeze (DEFRA 2011). This should be done as soon as possible, since compensatory sites can take some time to become suitable for their target species (Gilbert *et al.* 2010) and may not achieve the same plant communities as natural sites (Mossman *et al.* 2012).
 7. The needs of species currently resident in PAs for which climate will become less suitable should be balanced against the needs of species of conservation concern that colonise these sites during range expansions. The optimal balance will depend on the location of each reserve within the UK, and the importance of the site to the species it protects on an international level. Many birds associated with upland and montane habitats in the UK are of international conservation importance (Pearce-Higgins *et al.* 2011), so care should be taken not to disadvantage these species through habitat management.
 8. The habitat requirements of species that might colonise new areas should be identified from the north of their current climatic range. This is because species often show variation in habitat use across their full geographic range (e.g. Suggitt *et al.* 2012) and the requirements towards the north of their current range are likely to be the most similar to areas that become climatically suitable.

Knowledge Gaps

The effectiveness of PAs in conserving biodiversity under climate change is an emerging field of study, and as such there are a large number of knowledge gaps that should be considered priorities for research.

- Likely changes in abundance within PAs in taxa other than birds are unknown, but if models were generated these could be compared to monitoring data to determine whether numbers observed are as expected by models.
- The impact of changes in habitat extent and quality on abundance is unknown for most species, along with the likely interactions of these impacts with climate change.
- There is a lack of population monitoring of most taxa, even birds listed on the Birds Directive (Pearce-Higgins *et al.* 2011), from non-PA land, which makes it difficult to quantify the effectiveness of PAs.
- The area requirements and habitat preferences of species that might colonise the UK are often unknown in their current ranges, and filling this knowledge gap would help inform future habitat creation in the UK.
- The likely effectiveness of PA management in retaining viable populations of species predicted to do badly under climate change is largely unknown, and the results of management actions should therefore be closely monitored.
- Little is understood about the genetic components of biodiversity, and how to protect genetic diversity using PAs (Gaston *et al.* 2006).

- Likely future changes in land use in the matrix surrounding PAs is difficult to predict, and how these changes will affect species' ability to colonise areas that become climatically suitable is therefore unknown.
- The ability of species to track climate change to colonise all newly suitable PAs is largely unknown, as dispersal distances are not well understood for most species (e.g. Jaeschke *et al.* 2012). This limits our ability to project the future utilisation of PAs by potential colonisers.

References

- Amar, A., Grant, M., Buchanan, G., Sim, I., Wilson, J., Pearce-Higgins, J. W. and Redpath, S. (2011) Exploring the relationship between wader declines and current land-use in the British uplands. *Bird Study* **58**, 13-26
- Anderson, B. A. and Ohlemüller, R. (2011) Climate Change and Protected Areas: How well do British Rare Bryophytes fare? pp. 409-426 in *Bryophyte Ecology and Climate Change* (eds Z. Tuba, N. G. Slack and L. R. Stark). Cambridge University Press.
- Araújo, M. B., Cabeza, M., Thuiller, W., Hannah, L. and Williams, P. H. (2004) Would climate change drive species out of reserves? An assessment of existing reserve-selection methods. *Global Change Biology*, **10**, 1618-1626
- Araújo, M. B., Alagador, D., Cabeza M., Nogués-Bravo, D. and Thuiller, W. (2011) Climate change threatens European conservation areas. *Ecology Letters*, **14**, 484-492
- Arroyo, B., Leckie, F., Amar, A., Hamilton, J., McCluskie, A. and Redpath, S. (2005) *Habitat use and range management on priority areas for hen harriers: 2004 report*. NERC/Centre for Ecology & Hydrology, 45pp. (CEH Project Number: C02018)
- Bagchi, R., Crosby, M., Huntley, B., Hole, D. G., Butchart, S. H. M., Collingham, Y., Kalra, M., Rajkumar, J., Rahmani, A., Pandey, M., Gurung, H., Trai, L. T., Van Quang, N. & Willis, S. G. (2013) Evaluating the effectiveness of conservation site networks under climate change: accounting for uncertainty. *Global Change Biology* **19**, 1236-1248
- Bate, R., Bee, E., Jarvis, D. and Devine-Wright, P. (2010) Is MPS1 meeting its landscape and nature conservation objective. *Mineral Planning* **131**, 5. Available at http://nora.nerc.ac.uk/15454/1/Article_for_MP_mag_v3.pdf last accessed 13/08/2012
- BRANCH partnership (2007) *Planning for biodiversity in a changing climate* – BRANCH project Final Report, Natural England, UK.
- Brereton, T., Roy, D.B., Middlebrook, I., Botham, M. and Warren, M. (2011) The development of butterfly indicators in the United Kingdom and assessments in 2010. *Journal of Insect Conservation* **15**, 139-151
- Brooker, R., Ahrends, A., Bond, C., Castellazi, M., Ellis, C., Gimona, A., Harding, A., Kirkbridge, V., Poggio, L., Rowan, J., Wignall R. (2014) Summary Report: Climate Change Risk-Based Assessment for Notifiable Features in Scotland.

Carroll, C., Dunk, J. R. and Moilanen, A. (2010) Optimizing resiliency of reserve networks to climate change: multispecies conservation planning in the Pacific Northwest, USA. *Global Change Biology* **16**, 891-904

Carroll, M. J., Dennis, P., Pearce-Higgins, J. W. and Thomas, C. D. (2011) Maintaining northern peatland ecosystems in a changing climate: effects of soil moisture, drainage and drain blocking on craneflies. *Global Change Biology* **17**, 2991-3001

Chape, S., Harrison, J, Spalding, M. and Lysenko, I. (2005) Measuring the extent and effectiveness of protected areas as an indicator for meeting global biodiversity targets. *Philosophical Transactions B.* **360**, 443-455

Christie, M. and Rayment, M. (2012) An economic assessment of the ecosystem service benefits derived from the SSSI biodiversity conservation policy in England and Wales. *Ecosystem Services* **1**, 70-84.

Davies, H., Brereton, T. M., Roy, D. B. and Fox, R. (2007) Government targets for protected area management: will threatened butterflies benefit? *Biodiversity Conservation* **16**, 3719-3736

DEFRA (2011a) *Developing tools to evaluate the consequences for biodiversity of options for coastal zone adaptation to climate change. A study modelling the risk of loss from flooding of lowland open-water and wetland priority BAP habitats in the coastal floodplain under a range of sea-level rise scenarios.* Entec UK Limited.

DEFRA (2011b) *Biodiversity 2020: A strategy for England's wildlife and ecosystem services.* DEFRA, London.

Dockerty, T., Lovett, A. and Watkinson, A. (2003) Climate change and nature reserves: examining the potential impacts, with examples from Great Britain. *Global Environmental Change* **13**, 125-135

Dodd, A., Williams, G., Bradbury, R., Hardiman, A., Lonergan, A. and Watts, O. (2012) Protected Areas and Wildlife in Changing Landscapes: The Law and Policy Context for NGO Responses to Climate Change in the UK. *Journal of International Wildlife Law & Policy* **15**, 1-24.

Dodd, A., Hardiman, A., Jennings, K. and Williams, G. (2010) Protected areas and climate change: Reflections from a practitioner's perspective. *Utrecht Law Review* **6**, 141-150.

Donald, P. F., Sanderson, F. J., Burfield, I. J., Bierman, S. M., Gregory, R. D. and Waliczky, Z. (2007) International conservation policy delivers benefits for birds in Europe. *Science* **317**, 810-813.

Eastwood, A., Nijnik, M., Brooker, R., Pakeman, R., Artz, R., Norton, L., Ross, L., Bullock, J., Vellinga, N., Albon, S., Fielding, D., Irvine, R.J., Ramsay, S. & Cooksley, S. (2013) *Nature conservation and ecosystem service delivery.* JNCC Report no. 492

- Franks, S. E., Pearce-Higgins, J. W., Ausden, M., Massimino, D. (in press) Increasing the Resilience of the UK's Special Protection Areas (SPAs) to Climate Change. *Report to Natural England*
- Fuller, R. A., McDonald-Madden, E., Wilson, K. A., Carwardine, J., Grantham, H. S., Watson, J. E. M., Klein, C. J., Green, D. C. and Possingham, H. P. (2010) Replacing underperforming protected areas achieves better conservation outcomes. *Nature* **466**, 365-367
- Game, E. T. Bode, M., McDonald-Madden, E., Grantham, H. S. and Possingham, H. P. (2009). Dynamic marine protected areas can improve the resilience of coral reef systems. *Ecology Letters* **12**, 1336-1346
- Gaston, K. J., Charman, K., Jackson, S. F., Armsworth, P. R., Bonn, A., Briers, R. A., Callaghan, C. S. Q., Catchpole, R., Hopkins, J., Kunin, W. E., Latham, J., Opdam, P., Stoneman, R., Stroud, D. A. and Tratt, R. (2006) The ecological effectiveness of protected areas: The United Kingdom. *Biological Conservation* **132**, 76-87
- Gilbert, G., Brown, A. F. and Wotton, S. R. (2010) Current dynamics and vulnerability to sea level rise of a threatened Bittern *Botarus stellaris* population. *Ibis* **152**, 580-589
- Gillingham, P. K., Bradbury, R. B., Roy, D. R., Anderson, B. J., Baxter, J. M., Bourn, N. A. D., Crick, H. Q. P., Findon, R. A., Fox, R., Franco, A., Hill, J. K., Hodgson, J. A., Holt, A. R., Morecroft, M. D., O'Hanlon, N. J., Oliver, T. H., Pearce-Higgins, J. W., Procter, D. A., Thomas, J. A., Walker, K. J., Walmsley, C. A., Wilson, R. J. and Thomas, C. D. (2015a) The effectiveness of protected areas in the conservation of species with changing geographic ranges. *Biological Journal of the Linnean Society* **115**, 707-717
- Gillingham, P. K., Alison, J., Roy, D. B., Fox, R., Thomas, C. D. (2015b) High abundances of species in protected areas in parts of their geographic distributions colonised during a recent period of climatic change. *Conservation Letters* **8**, 97-106
- Harrop, S. R. (2011) 'Living In Harmony With Nature'? Outcomes of the 2010 Nagoya Conference of the Convention on Biological Diversity. *Journal of Environmental Law*, **23**, 117-128
- Haskoning, R. (2006) *Coastal squeeze, saltmarsh loss and Special Protection Areas*. English Nature Research Report No 710. Peterborough: English Nature.
- Heller, N. E. and Zavaleta, E. S. (2009) Biodiversity management in the face of climate change: A review of 22 years of recommendations. *Biological Conservation* **142**, 14-32
- Hickling, R., Roy, D. B., Hill, J. K., Fox, R. and Thomas, C. D. (2006) The distribution of a wide range of taxonomic groups are spreading polewards. *Global Change Biology* **12**, 450-455
- Hiley, J. R., Bradbury, R. B., Holling, M., Thomas, C. D. (2013) Protected areas act as establishment centres for species colonizing the UK. *Proceedings of the Royal Society, B* **280**: 20122310

Hiley, J. R., Bradbury, R. B., Thomas, C. D. (2014) Introduced and natural colonists show contrasting patterns of protected area association in UK wetlands. *Diversity and Distributions* **20**: 943-951.

HM Government (2013) *The National Adaptation Programme Making the country resilient to a changing climate*. The Stationary Office, London

Hodgson, J. A., Thomas, C. D., Dytham, C., Travis, J. M. J., & Cornell, S. J. (2012) The speed of range shifts in fragmented landscapes. *PLoS ONE* **7**: e47141.

Hodgson, J. A., Thomas, C. D., Cinderby, S., Cambridge, H., Evans, P. and Hill, J. K. (2011) Habitat re-creation strategies for promoting adaptation of species to climate change. *Conservation Letters* **4**, 298-297

Hodgson, J. A., Thomas, C. D., Wintle, B. and Moilanen, A. (2009) Climate change, connectivity and conservation decision making: back to basics. *Journal of Applied Ecology* **46**, 964-969

Hole, D. G., Huntley, B., Arinaitwe, J., Butchart, S. H. M., Collingham, Y. C., Fishpool, L. D. C., Pain, D. J. and Willis, S. G. (2011) Toward a management framework for networks of protected areas in the face of climate change. *Conservation Biology* **25**, 305-315

Hole, D. G., Willis, S. G., Pain, D. J., Fishpool, L. D., Butchart, S. H., Collingham, Y. C., Rahbek, C. and Huntley, B. (2009) Projected impacts of climate change on a continent-wide protected area network. *Ecology Letters* **12**, 420-431

Hopkins, J. J., Allison, H. M., Walmsley, C. A., Gaywood, M. and Thurgate, G. (2007) *Conserving biodiversity in a changing climate: guidance on building capacity to adapt*. DEFRA, London, UK.

Huntley, B., Hole, D. G. and Willis, S. G. (2012) Assessing the effectiveness of a protected area network in the face of climatic change. P345-364 in *Climate Change, Ecology and Systematics*, ed. Trevor R. Hodgkinson, Michael B. Jones, Stephen Waldren and John A. N. Parnell. Cambridge University Press

Huntley, B., Collingham, Y., Willis, S. and Hole, D. (2010) Protected Areas and Climatic Change in Europe: Introduction. In Natura 2000 and Climate Change – a Challenge. Götz Ellwanger, Axel Ssymank & Cornelia Paulsch Federal Agency for Nature Conservation. **Heft 118**: 7-27.

IPCC (2007) *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson, Eds., Cambridge University Press, Cambridge, UK

IPCC (2014) *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Barros, V.R., C.B. Field, D.J. Dokken, M.D. Mastrandrea, K.J. Mach, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R.

Mastrandrea, and L.L. White (eds.)). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. Chapter 23: Europe.

Jackson, S. F., Walker, K. and Gaston, K. J. (2009) Relationship between distributions of threatened plants and protected areas in Britain. *Biological Conservation* **142**, 1515-1522

Jaeschke, A., Bittner, T., Reineking, B. and Beierkuhnlein, C. (2012) Can they keep up with climate change? – Integrating specific dispersal abilities of protected Odonata in species distribution modelling. *Insect Conservation and Diversity* early online doi: 10.1111/j.1752-4598.2012.00194.x

Koleček, J., Schleuning, M., Burfield, I. J., Báldi, A., Böhning-Gaese, K., Devictor, V., Fernández-García, J. M., Hořák, D., Van Trunhout, C. A. M., Hnatyna, O. & Reif, J. (2014) Birds protected by national legislation show improved population trends in Eastern Europe. *Biological Conservation* **172**, 109-116

Kostyack, J., Lawler, J. J., Goble, D. D., Olden, J. D and Scott, M. (2011) Beyond Reserves and Corridors: Policy Solutions to Facilitate the Movement of Plants and Animals in a Changing Climate. *Bioscience* **61**, 713-719

Kujala, H, Araújo, M. B., Thuiller, W. and Cabeza, M. (2011) Misleading results from conventional gap analysis – Messages from the warming north. *Biological Conservation* **144**, 2450-2458

Lawler, J. J. and Hepinstall-Cymerman, J. (2010) Conservation planning in a changing climate: Assessing the impacts of potential range shifts on a reserve network. Ch 15 in S.C. Trombulak and R.F. Baldwin (eds.), *Landscape-scale Conservation Planning*, Springer

Lawson, C. R., Bennie, J. J., Thomas, C. D., Hodgson, J. A. & Wilson, R. J. (2014) Active management of protected areas enhances metapopulation expansion under climate change. *Conservation Letters* **7**, 111-118.

Lawton, J.H., Brotherton, P.N.M., Brown, V.K., Elphick, C., Fitter, A.H., Forshaw, J., Haddow, R.W., Hilborne, S., Leafe, R.N., Mace, G.M., Southgate, M.P., Sutherland, W.J., Tew, T.E., Varley, J. and Wynne, G.R. (2010). Making Space for Nature: a review of England's wildlife sites and ecological network. Report to Defra.

Lemieux, C. J. and Scott, D. J. (2005) Climate change, biodiversity conservation and protected area planning in Canada. *The Canadian Geographer* **49**, 384-399

Loarie, S.R., Duffy, P. B., Hamilton, H., Asner, G. P., Field, C. B. and Ackerly, D. D. (2009) The velocity of climate change. *Nature* **462**, 24-31

MacGregor, N. A. & van Dijk, N. (2014) Adaptation in Practice: How Managers of Nature Conservation Areas in Eastern England are Responding to Climate Change. *Environmental Management* **54**: 700-719

Mascia, M and Pailler, S. (2011) Protected area downgrading, downsizing and degazettement (PADDD) and its conservation implications. *Conservation Letters* **4**, 9-20

Mitchell, R. J., Rose, R. J. and Palmer, S. C. F. (2009) The effect of restoration techniques on non-target species: case studies in moorland ecosystems. *Applied Vegetation Science* **12**, 91-91

Monzón, J., Moyer-Horner, L. and Palamar, M. B. (2011) Climate Change and Species Range Dynamics in Protected Areas. *Bioscience* **61**, 752-761

Mossman, H. L., Davy, A. J. & Grant, A. (2012) Does managed coastal realignment create saltmarshes with 'equivalent biological characteristics' to natural reference sites? *Journal of Applied Ecology* early online doi: 10.1111/j.1365-2664.2012.02198.x

Natural England (2009) *Guidance on dealing with the changing distribution of tree species*. Natural England Technical Information Note TIN053. Natural England, Sheffield.

Natural England (2012). *Natural England's climate change risk assessment and adaptation plan*. Natural England General Publication, Number 318.

Natural England and RSPB (2014). *Climate Change Adaptation Manual: evidence to support nature conservation in a changing climate*. Natural England Publication NE 546. Natural England, Sheffield.

Natural England (2015) Improvement Programme for England's Natura 2000 Sites Climate change theme plan: Developing a strategic approach to climate change adaptation. Natural England, York.

Newson, S. E., Oliver, T. H., Gillings, S., Crick, H. Q. P., Morecroft, M. D., Duffield, S. J., Macgregor, N. J. & Pearce-Higgins, J. W. (2014) Can site and landscape-scale environmental attributes buffer bird populations against weather events? *Ecography* **37**, 872-882.

Normand, S., Svenning, J-C. and Skov, F. (2007) National and European perspectives on climate change sensitivity of the habitats directive characteristic plant species. *Journal for Nature Conservation* **15**, 41-53

Oliver, T., Stefanescu, C., Páramo, F., Brereton, T. & Roy, D. B. (2014) Latitudinal gradients in butterfly population variability are influenced by landscape heterogeneity. *Ecography* **37**, 863-871

Oliver, T. H., Thomas, C. D., Hill, J. K., Brereton, T. and Roy, D. B. (2012) Habitat associations of thermophilous butterflies are reduced despite climatic warming. *Global Change Biology* **18**, 2720-2729

Oliver, T. H., Hill, J. K., Thomas, C. D., Brereton, T. and Roy, D. B. (2009) Changes in habitat specificity of species at their climatic range boundaries. *Ecology Letters* **12**, 1091-1102

Opdam, P. and Wascher, D. (2004) Climate change meets habitat fragmentation: linking landscape and biogeographical scale levels in research and conservation. *Biological Conservation* **117**, 285-297

Pateman, R., Hill, J. K., Roy, D. B., Fox, R. and Thomas, C. D. (2012) Temperature-dependent alterations in host use drive rapid range expansions. *Science* **336**, 1028-1030

Pearce-Higgins, J. W. (2011a) Modelling conservation management options for a southern range-margin population of Golden Plover *Pluvialis apricaria* vulnerable to climate change. *Ibis* **153**, 345-356

Pearce-Higgins, J. W. (2011b). How Ecological Science Can Help Manage the Effects of Climate Change: A Case Study of Upland Birds pp. 397-414 in *The Changing Nature of Scotland*, eds. S.J. Marrs, S. Foster, C. Hendrie, E.C. Mackey, D.B.A. Thompson. TSO Scotland, Edinburgh

Pearce-Higgins, J. W., Johnston, A., Ausden, M., Dodd, A., Newson, S. E., Ockendon, N., Thaxter, C. B., Bradbury, R. B., Chamberlain, D. E., Jiguet, F., Rehfisch, M. M., and Thomas, C. D. (2011) *Final Report to the Climate Change Impacts on Avian Interests of Protected Area Networks (CHAINSPAN) Steering Group*. Available at <http://randd.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&Completed=2&ProjectID=16731> last accessed 10/08/2012

Pearce-Higgins, J. W., Dennis, P., Whittingham, M. J. and Yalden, D. W. (2010) Impacts of climate on prey abundance account for fluctuations in a population of a northern wader at the southern edge of its range. *Global Change Biology* **16**, 12-23

Pearson, R. G. and Dawson, T. P. (2005) Long-distance plant dispersal and habitat fragmentation: identifying conservation targets for spatial landscape planning under climate change. *Biological Conservation* **123**, 389-401

Pellatt, M. G., Goring, S. J., Bodtke, K. M. and Cannon, A. J. (2012) Using a Down-Scaled Bioclimate Envelope Model to Determine Long-Term Temporal Connectivity of Garry oak (*Quercus garryana*) Habitat in Western North America: Implications for Protected Area Planning. *Environmental Management* **49**, 802-815

Peters, R. L. and Darling, J. D. S. (1985) The greenhouse effect and nature reserves. *Bioscience* **35**, 707-717

Pyke, C. R. and Marty, J. (2005) Cattle grazing mediates climate change impacts on ephemeral wetlands. *Conservation Biology* **19**, 1619-1625

Pyke, C. R. and Fischer, D. T. (2005) Selection of bioclimatically representative biological reserve systems under climate change. *Biological Conservation* **121**, 429-441

Rayfield, B., James, P. M. A., Fall, A. and Fortin, M-J (2008) Comparing static versus dynamic protected areas in the Québec boreal forest. *Biological Conservation* **141**, 438-449

RSPB (2010) *Futurescapes: Space for nature, land for life*. Available at http://www.rspb.org.uk/Images/futurescapesuk_tcm9-253866.pdf last accessed 28/08/2012

RSPB (2007) *Climate change: Wildlife and adaptation. 20 Tough questions, 20 rough answers*. Available at

http://www.rspb.org.uk/Images/climatechange20questions_tcm9-170121.pdf last accessed 28/08/2012

Schweiger, O., Heikkinen, R. K., Harpke, A., Hickler, T., Klotz, S., Kudrna, O., Kühn, I., Pöyry, J. and Settele, J. (2012) Increasing range mismatching of interacting species under global change is related to their ecological characteristics. *Global Ecology and Biogeography* **21**, 88-99

Singh, N. J. and Milner-Gulland, E. J. (2011) Conserving a moving target: planning protection for a migratory species as its distribution changes. *Journal of Applied Ecology* **48**, 35-46

Smithers, R. J., Cowan, C., Harley, M., Hopkins, J. J., Pontier, H., Watts, O. (2008) England Biodiversity Strategy Climate Change Adaptation Principles Conserving biodiversity in a changing climate. Department for Environment, Food and Rural Affairs, London, UK.

Soutullo, A. (2010) Extent of the global network of terrestrial protected areas. *Conservation Biology* **24**, 362-363

Stein, A., Gerstner, K. & Kreft, H. (2014) Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales. *Ecology Letters* **17**, 866-880

Strange, N., Thorsen, B. J., Bladt, J., Wilson, K. A. and Rahbek, C. (2011) Conservation policies and planning under climate change. *Biological Conservation* **144**, 2968-2977

Suggitt, A. J., Stefanescu, C., Páramo, F., Oliver, T., Anderson, B. J., Hill, J. K., Roy, D. B., Brereton, T. and Thomas, C. D. (2012) Habitat associations of species show consistent but weak responses to climate. *Biology Letters* **8**, 590-593

Thomas, C. D., Gillingham, P. K., Bradbury, R. B., Roy, D. B., Anderson, B. J., Baxter, J. M., Bourn, N. A. D., Crick, H. Q. P., Findon, R. A., Fox, R., Hodgson, J. A., Holt, A. R., Morecroft, M. D., O'Hanlon, N. J., Oliver, T. H., Pearce-Higgins, J. W., Procter, D. A., Thomas, J. A., Walker, K. J., Walmsley, C. A., Wilson, R. J. and Hill, J. K. (2012) Protected Areas facilitate species range expansions *Proceedings of the National Academy of Sciences USA* **109**, 14063-14068

Trivedi, M. R., Morecroft, M. D., Berry, P. M. and Dawson, T. P. (2008) Potential effects of climate change on plant communities in three montane nature reserves in Scotland, UK. *Biological Conservation* **141**, 1665-1675

Verschuuren, J. (2010) Climate Change: Rethinking Restoration in the European Union's Birds and Habitats Directives. *Ecological Restoration* **28**, 431-439.

Virkkala, R., Pöyry, J., Heikkinen, R. K., Lehikoinen, A. & Valkama, J. (2014) Protected areas alleviate climate change effects on northern bird species of conservation concern. *Ecology and Evolution* early online, doi: 10.1002/ece3.1162

Virkkala, R., Heikkinen, R. K., Fronzek, S., & Leikola, N. (2013) Climate Change, Northern Birds of Conservation Concern and Matching the Hotspots of Habitat Suitability with the Reserve Network. *PLoS One* **8**(5): e63376

Virkkala, R. and Rajasärkkä, A. (2011) Climate change affects populations of northern birds in boreal protected areas. *Biology Letters* **7**, 395-398

Virkkala, R., Marmion, M., Heikkinen, R. K., Thuiller, W. and Luoto, M. (2010) Predicting range shifts of northern bird species: Influence of modelling technique and topography. *Acta Oecologica* **36** (2010) 269-281

Williams, J. M., ed. (2006) *Common Standards Monitoring for Designated Sites: First Six Year Report. Summary*. Peterborough, JNCC.

Willis, S. G., Hill, J. K., Thomas, C. D., Roy, D. B., Fox, R., Blakeley, D. S. and Huntley, B. (2009) Assisted colonisation in a changing climate: a test-study using two U. K. butterflies. *Conservation Letters* **2**, 45-51

Wilson, L., Astbury, S., Bhogal, A. and McCall, R. (2010). Climate Vulnerability Assessment of Designated Sites in Wales. CCW Science Report No. 942. Countryside Council for Wales, Bangor.

WWF (2004) *How effective are protected areas? A preliminary analysis of forest protected areas by WWF – the largest ever global assessment of protected area management effectiveness*. Available from <http://www.panda.org/downloads/forests/protectedareamanagementreport.pdf> last accessed 20/08/2012